

Evaluation of the Phytochemical and Antioxidant Properties of Cold, Hot Water and Wine Extracts Produced from *Ficus capensis* LeafUche Dennis-Eboh^{1*}, Fidelis I. Achuba², Betty O. George²¹Department of Medical Biochemistry, Faculty of Basic Medical Sciences, Delta State University, Abraka, Delta State, Nigeria²Department of Biochemistry, Faculty of Science, Delta State University, Abraka, Delta State, Nigeria**ABSTRACT**

Red wine consumption has increased tremendously due to the apparent health benefits and pleasure derived from red wines. In the quest to search for alternative sources of raw material for wine production, the phytochemical and antioxidant properties of the cold and hot water extracts of *Ficus capensis* leaf as well as the wine produced from hot water extract of *F. capensis* leaf were evaluated. The qualitative screening of phytochemical constituents in cold and hot water extracts of *F. capensis* leaf revealed the presence of flavonoids, tannins, phenols, steroids, carbohydrates, proteins, amino acids, alkaloids and phytate. The quantitative phytochemical screening of the cold and hot water extracts as well as the wine extracts of *F. capensis* using HPLC method revealed the presence of important bioactive ingredients with the concentration of the phenolic and flavonoid compounds of *F. capensis* wine significantly ($p < 0.05$) higher than the cold and hot water extracts. The wine extract exhibited significantly ($p < 0.05$) decrease in the antioxidant activity against DPPH, nitric oxide, total antioxidant capacity and FRAP compared to the cold and hot water extracts. Moderate consumption of *F. capensis* wine would help to protect the organism against oxidative damage to lipids, proteins, and nucleic acids caused by the phenols in the drink. Moderate consumption of *F. capensis* wine would help to protect the organism against oxidative damage to nucleic acids lipids, and proteins caused by the phenols in the drink.

Keywords: *Ficus capensis*, DPPH, Phytochemicals, Hot water extract, Cold water extract, wines, Antioxidants

Received 30 March 2023

Revised 10 April 2023

Accepted 12 April 2023

Published online 01 May 2023

Copyright: © 2023 Dennis-Eboh *et al.* This is an open-access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.**Introduction**

Wines have been made from fruits, herbs, vegetables, and sometimes a combination of both fruits and vegetables for decades.^{1,2} Making wine from vegetables has enhanced people's exposure to the plant's beneficial nutrients, chemicals, and medicinal properties.³ Most vegetables lack the necessary nutrients for yeast feeding, and water to form a stable drink naturally hence must is altered for local vegetable fermentation to produce their wines. Veggies with low quantity of fermentable sugars must be "chaptalized" in order for the resulting wine to have acceptable alcohol levels.² As a result, sucrose is added as the primary carbohydrate so that fruits with high acid levels (typically citric or malic acid) can hydrolyze sucrose into fermentable fructose and glucose sugars.¹ Some fruits and veggies have naturally high levels of acidity, making them unsuitable for making a tasty fruit wine. Therefore, the "must" is typically diluted with water to reduce the acidity to a more tolerable level prior to fermentation.⁴ These wines produced from fruits and vegetables are named after the parent plant.⁵ Fermentation is a bioengineering that involves feeding sugars and nutrients in solution to a required microorganisms, which reciprocates by producing carbon dioxide gas and alcohol.⁶

*Corresponding author. E mail: duche@delsu.edu.ng
Tel: 08050375789

Citation: Dennis-Eboh U, Achuba FI, George BO. Evaluation of the Phytochemical and Antioxidant Properties of Cold, Hot Water and Wine Extracts Produced from *Ficus capensis* leaf. Trop J Phytochem Pharm. Sci. 2023; 2(2):48-54. <http://www.doi.org/10.26538/tjpps/v2i2.1>

Official Journal of Natural Product Research Group, Faculty of Pharmacy, University of Benin, Benin City, Nigeria.

As a result, fruits rich in acid (usually citric or malic acid) are added to the mix as the major carbohydrate, and the sucrose they contain is hydrolyzed into fermentable fructose and glucose sugars.¹ Some fruits and veggies have naturally high levels of acidity, making them unsuitable for making a tasty fruit wine. Therefore, the "must" is typically diluted with water to reduce the acidity to a more tolerable level prior to fermentation. At the moment, most of the wine production processes are relying on *Saccharomyces cerevisiae* strains that allow rapid and reliable fermentations, reduce the risk of sluggish or stuck fermentations and prevent microbial contaminations.⁷ Yeast starter cultures that are deliberately chosen for the winemaking process based on scientifically proven features often complement and optimize the raw material quality and individual characteristics⁴ of the alcohol, resulting in a more desirable outcome.^{7,8} These musts and wines are fermented and aged, and their alcohol concentration typically ranges from 5 to 13%.⁹

In Nigeria, red wine consumption has increased tremendously due to the apparent health benefits and pleasure derived from red wines.¹⁰ European countries such as France, Spain, and Italy contribute for around 60% of overall imports to Nigeria, while the Republic of South Africa and America account for approximately 22% and 8% of the nation's total imports, respectively.^{11,12} The rise in import costs for good-quality red wine, the rapid depreciation of the Nigerian currency, and competition for the use of these fruits between wineries and man as a result of wines being produced from veggies and fruit due to grape scarcity in the world's tropical regions today have necessitated the search for plants with physicochemical and antioxidant attributes for wine production.¹³ *Ficus capensis*, often known as *Ficus sur*, is a member of the Moraceae family. The plant matures into a deciduous tree that can reach a height of 35 to 40 meters. It has extensive roots, branches, and large green leaves (115-131ft). Young leaves have a glabrous top surface and a hairy below surface.¹⁴ They are spirally and alternately organized, with uneven margins and elliptical to ovate

forms. All year long, *F. capensis* bears fleshy fruits along the trunk and main branches, either in a single or branched raceme.¹⁵ It is a well-known medicinal plant that has long been used to treat a variety of diseases and disorders, such as diabetes, liver disorders, diarrhea, inflammatory conditions, hemorrhoids, respiratory ailments, and urinary diseases. The fruits of *F. capensis* are used to cure dry cough, loss of voice, kidney and spleen disease, intestine astringency, leucorrhoea, a burning feeling, weariness, and urinary tract infection.¹⁶ The fruit extract has antiulcer qualities, and the component -sitosterol present in the fruit has substantial hypoglycemic activity, which means it lowers blood glucose levels.¹⁷ Ayurvedic practitioners employ the fruits of *F. capensis* to treat bleeding conditions such as nose hemorrhage, menorrhagia, and rectal bleeding.¹⁸ The leaves and roots have been shown in Sudan and Nigeria to exhibit biologically beneficial drug-like qualities such as microbial inhibitory activity,¹⁹ antiulcer activity,²⁰ antisickling activity,²¹ antidiarrheal activity,²² leukocyte mobilization, and immune-boosting action.^{23,14} Phytochemical characteristics of *F. capensis* leaf, root, and seed extracts have previously been documented in the literature.^{24,14,16} There has been no research on the phytochemical qualities of hot water and wine extracts created from *F. capensis* leaf, thus the current work explores the phytochemical and antioxidant properties of cold, hot, and wine extracts produced from *F. capensis* leaf.

Materials and Method

Fresh matured *F. capensis* leaf separated from stalks was collected from a local bush in 2019 in Abraka, Ethiopia East Local Government Area of Delta State. It was identified and authenticated with a voucher specimen (UBH-f331) by a taxonomist Dr. H.A. Akinnibosun, at the Division of Botany, Faculty of Life Sciences, University of Benin, Nigeria.

Preparation of *F. capensis* extract/sample

Cold water sample

The plant tissue homogenization technique was applied for the preparation of cold water leaf samples of *F. capensis*. About 100 g *F. capensis* leaf was weighed, and crushed to a smooth paste using mortar and pestle. Before filtering, about one liter of deionized water was added and shaken vigorously at 36.8°C for 5–10 minutes. The centrifugation of filtrate was done for sample clarification.²⁵

Hot water sample

The decoction technique was applied for the preparation of the hot water leaf sample of *F. capensis*. This was achieved by heating 150g of the fresh leaf in 1 litre of water for 15 - 20 minutes at 100 degrees Celsius and thereafter left to cool down.²⁵ Both sample extracts were appropriately labeled and stored in a refrigerator at 4 °C to avoid deterioration and wastage for future use.

Bioactive compounds of cold and hot water extracts of *F. capensis* determination

The qualitative phytochemical screening was done using standard principles to identify alkaloids,²⁶ amino acids, carbohydrates, phenolic compounds and tannins, phytosterol, cardiac glycoside, proteins (millon's test),²⁵ anthraquinones, anthocyanosides, flavonoids and saponins.²⁷

Chromatographic analysis of phytochemical compounds in the cold, hot and wine extracts of *F. capensis*

The mobile phase is made up of 1% aqueous acetic acid solution (Solvent A) and acetonitrile (Solvent B), with a flow rate of 2ml/min. The sample injection volume was 5l, and the column temperature was set to 28 °C. To achieve gradient elution, the percentage of solvent B to solvent A was changed. The mobile phase composition was returned to its initial solvent B: solvent A: 10: 90 ratio in 55 minutes, and it was left to stand for another 10 minutes before another sample injection. The entire analysis time per sample was 65 minutes. The HPLC chromatograms were detected using a photo diode array UV detector at three different wavelengths: 272, 280, and 310 nm, depending on the chemicals investigated. Compounds were identified using retention

time and spiking with standards under the same conditions. The integrated peak area was measured and the content was estimated to complete the sample measurement.

Calculation

Total and individual parameter specifications

The following are the total and individual standard concentrations of cold, hot, and clarified wine:

Parameter concentration (mg/g) = (Peak area)/(Standard peak area) x Standard concentration

Total phenols quantitative estimation²⁸ reducing sugar determination assay,²⁹ total flavonoids estimation,³⁰ reducing sugar determination assay,²⁹ alkaloids,³¹ and phytate.³²

Assay for antioxidants

The antioxidant activity and free radical scavenging of *F. capensis* cold, hot, and wine extracts were determined in vitro using the 1, 1-Diphenyl-1-picrylhydrazyl (DPPH) assay,³³ Nitric Oxide (NO),³⁴ FRAP Assay³⁵, and Total Antioxidant Capacity.³⁶

Statistical investigation

Using SPSS version 22.0, the generated data were subjected to the student T test. At p < 0.05, the mean differences were judged significant.



Results and Discussion

Table 1 shows the qualitative phytochemical composition of cold and hot water extracts of *F. capensis* leaf. The presence of significant flavonoids, tannins, phenols, steroids, carotenoids, terpenes, alkaloids, and phytate was discovered in the cold and hot water extracts of *F. capensis* leaf (Table 1), (Figure 1), and (Figure 2). The study revealed that the hot water and wine extracts were high in phenol, flavonoids, reducing sugar, and alkaloid content. (Table 2). The wine had more phenol, alkaloids, and reducing sugars than flavonoid and phytate (Table 2). Flavonoids are water-soluble chemicals that diminish free radicals by quenching, up-regulating, or protecting antioxidant defenses, as well as chelating radical intermediate molecules.³⁷ The HPLC phytochemical constituents of cold and hot water extracts of *F. capensis* leaf (Figure 1- 4) was found to contain Quercetin > Kaemferol > Geranyl acetone > Rutin > Alpha Caryophyll > Linalool > Orientin > Catechin > Epigallocatechin > Myricetin > Scytaline > Lupeol > Cetylbehenite > Luteolin, which are very important bioactive phytochemicals. In the cold water extract of *F. capensis*, the highest quantities were Quercetin (96.25 ppm) and Kaemferol (31.74 ppm), while the lowest values were Luteolin (0.61 ppm) and Cetylbehenite (0.46 ppm). In the hot water extract of *Ficus capensis*, the greatest concentrations were Quercetin (98.44 ppm) and Kaemferol (34.4 ppm), while the lowest values were Cetylbehenite (0.64 ppm) and Luteolin (0.61 ppm). The hot water extract's total phytochemical concentrations (21.25 ppm) were found to be greater than the cold water extract's total phytochemical concentrations (19.39 ppm). As a result, the presence of these essential phytochemicals in the water extracts of *F. capensis* highlights the leaf's potential nutritional and therapeutic value. Phytochemicals are bioactive substances found naturally in plants that have protective and disease-preventive characteristics, making them useful in medicine and pharmacology.^{38,39,40,41,42} Alkaloids protect plants against herbivores, play metabolic roles in living organisms, and have pharmacological activities such as analgesic, antimicrobial, antibacterial, and antibiotic properties. Plant flavonoids and phenols have been shown to have antioxidant, free radical scavenging, anti-inflammatory, and anticarcinogenic effects.^{40,41,42} plant-derived phenol molecules have been shown to contribute to the color, sensory, and antioxidant qualities of food.^{42,43,44,45} Several methods have been proposed to explain the antioxidant activity of phenolic compounds, including reducing capacity, chain initiation prevention, binding of transition metal ion catalysts, peroxide breakdown, continuing hydrogen abstraction prevention, and radical scavenging.^{46,47,48,49} The findings confirm the work of^{50,51,52,16} who discovered that *F. capensis* has high nutritional value and can be used as a nutraceutical as well as a possible source of natural antioxidants that may slow the progression of various oxidative-related disorders.

Table 1: Qualitative Phytochemical Screening

Phytochemicals	Cold sample	Hot sample
Saponins	-	-
Phlobatannins	-	-
Cardiac glycosides	-	-
Flavonoids	+	+
Tannins	+	+
Phenols	+	+
Steroids	+	+
Terpenes	-	-
Carbohydrates	+	+
Protein	+	+
Amino acid	+	+
Gum and mucilage	-	-
Alkaloid	+	+
Phytate	+	+

Key: +Present, -Absent

**Figure 1:** Phytochemical profile of cold and hot water leaf extract of *F. capensis* with concentration CS – Cold sample; HS- Hot sample**Figure 2:** Phytochemical profile of cold and hot water leaf extract of *F. capensis* with concentration CS – Cold sample; HS- Hot sample

The work is also consistent with^{21,19,51,20}. Saponin was shown lacking in both extracts, as demonstrated by Oitoku's research.⁵³ In the ethanol extract of *F. capensis* leaves, Ezeigwe¹⁶ found a significant percentage of flavonoids, alkaloids, tannin, saponin, phate, oxalate, phenol, haemagglutinin, and cardiac glycosides. Wilfred⁵⁴ reported the presence of these phytochemicals in abundance in *F. capensis* leaf extracts, whereas Chizoruo⁵⁵ found the presence of varied levels of flavonoids, terpenes, and steroids in *F. capensis* leaf extracts. Saponins, cardiac glycosides, flavonoids, and tannins were found in the aqueous extracts of *F. capensis* thumb by Ayinde and Owolabi²², but alkaloids, cyanogenic glycosides, and anthracene derivatives were not. The presence of phytochemical elements may differ depending on the climatic and soil makeup of the place where the plant was harvested.^{55,56} Figure 5 - 6 depicts the phytochemical profile and HPLC chromatogram of *F. capensis* wine, which included more phytochemicals in the following order of concentration: Resveratrol, quercetin, trans-E-viniferin, gallic acid, chlorogenic acid, beta carotene, delphinidin, caffeic acid, lutein, epicatechin, petunidin, cyanidin, stigmasterol, catechin, beta sitosterol, and taxifolin are all examples of phytonutrients. Resveratrol (11.96 ppm) had the highest concentration in the wine, whereas taxifolin (0.11 ppm) had the lowest. The overall concentration of phytochemicals is 29.69 ppm. The phenolic chemicals kaempferol, quercetin, and catechin are capable of chelating metallic ions.^{46,47,48,49,52,53,54} These secondary bioactive metabolites play a significant function in red wine by acting as anti-alcohol agents.⁴⁶ The wine's higher phenolic content could be linked to the fermentation process's ability to convert conjugated forms of phenolic compounds to their free forms. This could possibly be due to the microbes producing proteolytic enzymes, which are responsible for hydrolyzing phenolic complexes into soluble - free phenols.⁵⁷

The most effective wine antioxidants include resveratrol, anthocyanins, and catechins.^{58,59} The levels of resveratrol, quercetin, and gallic acid in the wine were higher than the levels of other phytochemicals. Resveratrol has been found to play a role in cardiovascular disease prevention.⁵⁹ Quercetin has been shown to have health advantages in a variety of chronic conditions, including neurological disorders, diabetes, cardiovascular disease, and cancer.^{60,61} It can also prevent the oxidation of glutathione (GSH), which aids in the prevention of neurotoxicity caused by oxidative stress.^{62,63} It has been discovered to protect the brain from cytotoxicity caused by H₂O₂.⁶³ Trans-E-viniferin reduces reactive oxygen species while increasing glutathione (GSH) levels.^{64,65} Gallic acid (3,4,5-trihydroxybenzoic acid) is a naturally occurring antioxidant that may protect healthy people by preventing apoptosis.^{47,50} It is widely acknowledged that chlorogenic acid (CGA) offers various health benefits as an anti-hypertension drug.⁶⁶ Chlorogenic and caffeic acids, two of the most prominent hydroxycinnamic acids, have also been linked to a reduction in oxidative and inflammatory stress. Beta-carotene is a provitamin A that can be transformed into vitamin A (retinol), which plays important roles in the control of physiological activities in animal bodies.⁶⁷ The discovered anthocyanin profiles are delphinidin > petunidin > cyanidin. Anthocyanin is a powerful antioxidant and anti-inflammatory substance that can reduce the risk of acquiring diseases such as atherosclerosis. It promotes longevity and reduces oxidative stress.^{68,69,70} Caffeic acid has numerous therapeutic effects, including antioxidant, immunomodulatory, antibacterial, neuroprotective, antianxiety, antiproliferative, and anti-inflammatory qualities.^{71,72} Caffeic acid and chlorogenic acid have been demonstrated to inhibit key enzymes associated with hypertension and to protect the heart from prooxidant-induced oxidative damage in rats.⁷³ Lutein is a carotenoid that has anti-oxidative and anti-inflammatory qualities that protect against alcohol-induced liver injury^{68,74,75} ameliorate or even prevent age-related macular disease, which is the primary cause of blindness and vision impairment. Steroids and phytosterols have been shown to increase nitrogen retention in osteoporosis and wasting disease mice.³⁸ As a rich source of bioactive components and powerful antioxidants, *F. capensis* wine will guard against alcohol damage while also providing various health benefits such as anti-diabetic, anti-aging, anti-microbial, and chronic disease protection. However, the amount drunk should be limited because excessive alcohol use can have negative economic, medical, and societal implications.^{44,59,69}

Table 2: Quantitative Phytochemical and reducing sugar screening of the cold, hot water and wine extracts of *F. capensis* leaves using spectrophotometer



Phytochemicals	Cold sample	Hot sample	Clarified wine
Phenols (mg/gGAE)	0.03 ± 0.000a	0.04 ± 0.000ac	0.042 ± 0.001c
Flavonoids (mg/gGAE)	0.15 ± 0.010b	0.45 ± 0.080ac	0.074 ± 0.004d
Alkaloid (mg/gCAE)	0.17 ± 0.010a	0.22 ± 0.020ac	0.472 ± 0.006d
Phytate (mg/g)	0.37 ± 0.010b	0.14 ± 0.020bc	0.050 ± 0.001d
Reducing Sugars (mg/gGluE)	0.36 ± 0.010b	1.42 ± 0.010ac	1.528 ± 0.016d

Values are means ± standard deviations of triplicate determinations. Values not sharing common superscripts on the same column differ significantly ($p < 0.05$). The mean differences were considered significant at $p < 0.05$. GAE = Gallic acid equivalent, CAE = Catechin equivalent, GluE = Glucose equivalent.

Table 3: Free radical and *In-vitro* antioxidant activity of cold, hot water and wine extract of *F. capensis* leaves

Antioxidant parameter	DPPH (%inhibition)	FRAP (%inhibition)	NO (%inhibition)	TAC (mg/gGAE)
Cold sample	79.02 ± 2.94 ^a	1.60 ± 0.02 ^b	75.33 ± 0.33 ^a	0.09 ± 0.00 ^b
Hot sample	87.1 ± 0.92 ^b	2.22 ± 0.03 ^a	77.6 ± 1.73 ^a	0.24 ± 0.00 ^a
Wine sample	30.03 ± 15.57 ^c	0.02 ± 0.01 ^c	55.41 ± 4.09 ^b	0.27 ± 0.01 ^a

Values are means ± standard deviation of triplicate determinations. Values not sharing common superscripts on the same column differ significantly ($p < 0.05$). DPPH = 1,1-diphenyl-2-picrylhydrazyl, FRAP = Ferric reducing antioxidant power, NO (%inhibition) = Nitric Oxide and TAC (mg/gGAE) = Total antioxidant capacity.

**Figure 3:** Chromatogram of phytochemical composition of cold water leaf extract of *Ficus capensis***Figure 4:** Chromatogram of phytochemical composition of hot water leaf extract of *F. capensis*

F. capensis hot and cold water extracts demonstrated antioxidant efficacy against DPPH, nitric oxide, total antioxidant capacity, and FRAP activities. (Table 3). The DPPH radical assay is a rapid, dependable, and reproducible method for evaluating antioxidant free radical scavenging activity. DPPH is a stable nitrogen-centered radical that readily accepts an electron or a hydrogen radical to form a stable diamagnetic molecule.^{64,76,77} The DPPH test is an important procedure in determining a substance's antioxidant capacity.^{60,77,78} The FRAP test is another frequently used approach for determining a phytochemical compound's ability to give electrons that decrease oxidized intermediates.⁷⁷ Doris and Rosemary⁷⁹ discovered that increasing the concentration of extract increased the antioxidant activity of *F. capensis* leaves. According to Seun *et al.*,⁴⁷ radical scavenging and Fe^{2+} -chelating capacities are two probable mechanisms through which *F. capensis* leaves extract could be used to treat erectile dysfunction. *In vitro* antioxidant activity of *F. capensis* was also reported by Ehimwenma and Osarieme.⁸⁰

The antioxidant activity of the wine extract against DPPH, nitric oxide, total antioxidant capacity, and FRAP was reduced. (Table 3). This could occur as a result of the wine's aging. Jan *et al.*⁸¹ discovered that the antioxidant activity of several fruit extracts decreases with age. The antioxidant capacity of wine is thought to be highly associated with the type of phenolic compounds found in wine. This demonstrates one of the most important roles of phenols in antioxidant activity.^{40,42,44,47,48,49} Antioxidant activity is now regarded as one of the most important features of red wines, and it is linked to the presence of polyphenols such as flavonoids, phenolic acids, stilbenes, coumarins, lignoids, and so on.^{44,45,49,73,82,83} Moderate consumption of *F. capensis* wine would help to neutralize free radicals, protecting the organism from oxidative damage to lipids, proteins, and nucleic acids caused by the phenols in the drink.^{51,81}

Conclusion

This finding implies that phytochemical compounds discovered in the cold water extract of *F. capensis* leaf were also found in the hot water extract. Furthermore, *F. capensis* wine prepared from hot water extract using *Saccharomyces cerevisiae* includes bioactive ingredients that, when consumed in moderation, will prevent cardiovascular disorders as well as act as a preventative agent against alcohol damage.




Figure 5: Phytochemical profile of *F. capensis* wine




Figure 6: Chromatogram of phytochemical composition of *F. capensis* wine

Conflict of Interest

The authors declare no conflict of interest.

Authors' Declaration

The authors hereby declare that the work presented in this article is original and that any liability for claims relating to the content of this article will be borne by them.

Acknowledgements

Mr Ohwokevwo Oghenyore Andy, Dr Okpogono Joel, Dr Apiamu Augustin, Dr Egbule Egoamaka Great, Dr Umuokoro Koghwo Emuesiri and Mr Ajoh Alfred all contributed to the accomplishment of this work.

References

- Ebana RUB, Edet UO, Anosike KI, Etok CA, Kanu TO. Nutritional analysis and wine production potentials of *Telfairia occidentalis* (fluted pumpkin) leaves and *Cucumis sativus* L. (cucumber) using Baker's and palm wine yeast strains. WNOFNS. 2019; 22:12-30.

- Kantiyok BD, Mohammed SSD, Yabaya A. Fruit wine produced from pawpaw (*Carica papaya*) and watermelon (*Citrullus vulgaris*) using *saccharomyces cerevisiae* isolated from selected fermented foods. Sci World J. 2021; 16(4):419-427.
- Waheed D, Sadiya AD. Herbal wine formulation from fruits and vegetables wastes and peels. IJEAST. 2021; 5(9):129-133.
- Yahia EM, Maldonado Celis ME, Svendsen M. The contribution of fruit and vegetable consumption to human health. Fruit and Vegetable Phytochemicals: Chem and Human Health, 2nd Edition. 2017; 11:1-52.
- Chigozie VU, Okezie MU, Ajaegbu EE, Okoye FB, Esimone CO. Bioactivities and HPLC analysis of secondary metabolites of a morphologically identified endophytic *Aspergillus* fungus isolated from *Mangifera indica*. Nat Prod Res. 2022; 36(22):5884-5888.
- Roffler R, Randolph TW, Miller DA, Blanch HW, Prausnitz JM. Extractive bioconversions with nonaqueous solvents. In Extractive bioconversions. 2021; 133-172.
- Mulani SA, Tamboli F A, More H N, Kolekar YS, Mali N. Formulation and evaluation of medicated guava (*Psidium guajava* L.) wine using *Saccharomyces crevices* Var. HAU 1. Indian J Pharm Pharmacol. 2021; 8(2):168-170.
- Nardi T. Microbial resources as a tool for enhancing sustainability in winemaking. Microorganisms. 2020; 8(4):507.
- Lopes ACA, Eda SH, Andrade RP, Amorim JC, Duarte WF. New alcoholic fermented beverages—potentials and challenges. Fermented Beverages. 2019; 577-603.
- Schaefer R, Olsen J, Thach L. Exploratory wine consumer behavior in a transitional market: The case of Poland. Wine Econ. 2018; 7(1):54-64.
- Nwachukwu C, Hieu MV, Chládková, H, Fadeyi O. Strategy implementation drivers in correlation with strategic performance. J Mark Manag. 2019; 17(1): 19-38.
- Ohana-Levi N, Netzer Y. Long-Term Trends of Global Wine Market. Agriculture, 2023; 13(1):224-248
- Ofoedu CE, Ofoedu EO, Chacha JS, Owuamanam CI, Efekalam IS, Awuchi CG. Comparative Evaluation of Physicochemical, Antioxidant, and Sensory Properties of Red Wine as Markers of Its Quality and Authenticity. Int J Food Sci. 2022; 2022.
- Kevwe, B. E., Samuel, O. A., Omolola, T. F and Oluyemisi, F. K. *Ficus capensis* Thumb. (Moraceae): Review of Its Ethnomedicinal Uses, Pharmacological Activities and Phytochemical Constituents. Arch Curr Res Int. 2018; 12(13):1-7
- Pawlos Z, Chandravanshi BS, Yohannes W, Embiale A. Levels of selected metals in *Ficus sur* Forssk fruit and soil of the plant grown in different parts of Ethiopia. SINET: Ethiop J Health Sci. 2021; 44(1):1-12.
- Ezeigwe OC, Okpala CO, Ogana J, Akonam NE, Aziagba BO, Nnadi NN, Anekwe OJ, Nwobodo VOG. Comparative Phytochemical and Nutritional Profiles of *F. capensis* and *Cnidioscolus aconitifolius* Leaves. IJRIAS. 2020; 5(1):16-21.
- Nawaz H, Waheed R, Nawaz M. Phytochemical composition, antioxidant potential, and medicinal significance of *Ficus*. Modern Fruit Industry. 2020; 1(20).
- Van Vuuren S, Frank L. Southern African medicinal plants used as blood purifiers. J ethnopharmacol. 2020; 249:112434.
- Adebayo-Tayo R, Odeniyi A. Phytochemicals screening and Microbial inhibitory. Activities of *Ficus capensis*. Afr J Biomed Res. 2012; 15:35-40
- Eluka P, Nwodo F, Akah P, Onyeto C. Anti-ulcerogenic and antioxidant properties of the aqueous leaf extract of *F. capensis* in Wistar albino rats. MRJMMS. 2015; 3(1): 022-026

21. Uzoekwe NM, Mohammed JJ. Phytochemical, proximate and Mineral contents of leaves and back of *Ficus capensis*. J Appl Sci Environ Manag. 2015; 19(4):633-637.
22. Ayinde BA, Owolabi OJ. Effect of the aqueous extract of *F. capensis* Thunb (Moraceae) leaf on gastrointestinal motility. J Pharmacogn and Phytotherapy. 2009;1(3):031-035
23. Daikwo OA, Tende JA, Okey S, Eze ED, Isa AS. The Effect of Aqueous Extract of Leaf of *Ficus capensis* Thunb (Moraceae) on *in vivo* Leukocyte Mobilization in Wistar Rats. Br J Pharmacol Toxicol. 2012; 3(3):110-114
24. Adaji MU, Dauda JA, Ameh EM, Ocheme GW, Aaron PE. Evaluation of Antioxidant Potency, Antimicrobial Activity and Phytochemical Screening of *Ficus capensis* Leaves (Cape Fig). Int J Glob Sustain. 2020; 6(3):8-8.
25. Sahira KB, Cathrine L. General techniques involved in phytochemical analysis. Int J Adv Res Chem Sci. 2015; 2(4):25-35.
26. Evans WC. Trease and Evans pharmacognosy. 1997. 14th edn., W.B Saunders, London, UK
27. Osuntokun OT, Oluwafoise BO. Phytochemical screening of ten Nigerian medicinal plants. Int J Multidiscip Res Dev. 2015; 2(4): 390-396.
28. Singleton, V. and Rossi, J. Colorimetry of Total Phenolic Compounds with Phosphomolybdic-Phosphotungstic Acid Reagents. Am J Enol Vitic. 1965; 16:144-158.
29. Miller GL. Use of Dinitrosalicylic Acid Reagent for Determination of Reducing Sugar. J Anal Chem. 1959; 31: 426-428.
30. Jia ZS, Tang MC, Wu JM. The Determination of Flavonoid Contents in Mulberry and Their Scavenging Effects on Superoxide Radicals. Food Chem. 1999; 64: 555-559.
31. Shamsa F, Hamidreza M, Rouhollah G. Mohammadreza V. Spectrophotometric determination of total alkaloids in some Iranian medicinal plants. Thai J Pharm Sci. 2008; 32:17-20
32. Wilcox JR, Premachandra, GS. Young K.A. and Raboy V. Isolation of high seed inorganic P, low-phytate soybean mutants. Crop Sci. 2000; 40:1601-1605
33. Manzocco, L, Anese M, Nicoli MC. Antioxidant properties of tea extracts as affected by processing. LWT. 1998; 31 (7-8):694-698.
34. Marcocci I, Marguire JJ, Droyy-lefaiz MT, Packer L. The nitric oxide scavenging properties of ginkgo biloba extract. *Biochem Biophys Res Commun*.1994; 201:748-755.
35. Benzie FF, Strain JJ. Ferric reducing/antioxidant power assay: Direct measure of total antioxidant activity of biological fluids and modified version for simultaneous measurement of total antioxidant power and ascorbic acid concentration. Meth Enzymol. 1996; 299:15-27
36. Prieto P, Pineda M, Aguilar M. Spectrophotometric quantitation of antioxidant capacity through the formation of a phosphomolybdenum complex: specific application to the determination of vitamin E. Anal Biochem.1999; 269(2):337-41.
37. Pawaskar SM, Sasangan KC. Preliminary phytochemical and invitro-antimicrobial analysis of *Annona squamosa linn.* leaf extract. J Pharm Sci Res. 2017; 9(5):618.
38. Shadrach I, Banji A, Adebayo O. Nutraceutical potential of ripe and unripe plantain peels: A comparative study. Chem Int. 2020; 6(2):83-90.
39. Singayina, BE Keswet, LA Asiya, L Osaji, MJ. Assesment of Nurritional Analysis of *Bombax buonopozense* Found in Adamawa and Taraba States, Nigeria (Red Flowered Silk Cotton Tree). *Sciences*. 2022; 11(1):7-11.
40. Ngozi P Okolie, Falodun A and Oluseyi Davids. Evaluation of the antioxidant activity of root extract of pepper fruit (*Dennetia tripetala*), and its potential for the inhibition of Lipid peroxidation. Afri J Trad Complem and Altern Med 2014; 11(3):221-227.
41. Egharevba E, Chukwuemeke-Nwani P, Eboh U, Okoye E, Bolanle IO, Oseghale IO, Imieje VO, Erharuyi O, Falodun A. Antioxidant and Hypoglycaemic Potentials of the Leaf Extracts of *Stachytarphyta jamaicensis* (Verbenaceae). Trop J Nat Prod Res. 2019; 3(5):170-174.
42. Jucá MM, Cysne Filho FMS, de Almeida JC, Mesquita DDS, Barriga JRDM, Dias KCF, Vasconcelos SMM. Flavonoids: biological activities and therapeutic potential. Nat Prod Res. 2020; 34(5):692-705.
43. Ajai N, Elavarasan S, Priya AK, Akash S, Annie TJ, Bhuvana G. Experimental study on partial replacement of cement by metakaolin and GGBS. Mater Today: Proc. 2021; 37:3527-3530.
44. Kumar SS, Arya M, Nagbhushan P, Giridhar P, Shetty NP, Yannam SK, Mahadevappa P. Evaluation of various drying methods on bioactives, ascorbic acid and antioxidant potentials of *Talinum triangulare L.* foliage. Plant Foods Hum Nutr. 2020; 75(2):283-291.
45. Okechukwu VU, Amaefule AI, Omokpariola DO, Umeh TC, Oze NR, Omokpariola PL. The Impact of Processing Procedures on the Proximate and Phytochemical Composition of Unripe *Musa paradisiaca* (Plantain). Eur J Sustain Dev Res. 2022; 6(3):188.
46. Kaur P, Purewal SS, Sandhu KS, Kaur M, Salar RK. Millets: A cereal grain with potent antioxidants and health benefits. J Food Meas Charact. 2019; 13(1):793-806.
47. Zahrani NAA, El-Shishtawy RM, Asiri AM. Recent developments of gallic acid derivatives and their hybrids in medicinal chemistry. A review. Eur J Med Chem. 2020; 7
48. Egbune EO, Ezedom T, Anigboro AA, Aganbi E, Amata AI, Tonukari NJ. Antioxidants and antigenotoxic properties of *Rhizopus oligosporus* fermented cassava (*Manihot esculenta Crantz*). *Afr J Biochem Res*. 2022; 16(3):39-46
49. Odion EE, Falodun A and Adelusai SA. Total flavonoid, Total Phenolic and antioxidant potential of root bark extract and fractions of from *Cola rostrata* (Sterculiaceae) K. Schum. University of Benin. J Sci Tech. 2013; 1(2): 38-42.
50. Oluwajuyitan TD, Ijarotimi OS, Fagbemi TN. Plantain-based dough meal: Nutritional property, antioxidant activity and dyslipidemia ameliorating potential in high-fat-induced rats. Food Frontiers. 2022; 3(3):489-504.
51. Seun AF, Oboh G, Oyeleye SI, Aline BA. Aqueous extract from *F. capensis* leaves inhibits key enzymes linked to erectile dysfunction and prevent oxidative stress in rats' penile tissue. NFSJ. 2016; 4:15-21.
52. Shodehinde SA, Adefegha SA, Oboh G, Oyeleye SI, Olasehinde TA, Nwanna EE, Adebayo BC. Boligon AA. Phenolic composition and evaluation of methanol and aqueous extracts of bitter melon (*Momordica charantia L*) leaves on Angiotensin-I-converting enzyme and some pro-oxidant – induced Lipid peroxidation invitro. J Evid Based Complementary Altern. 2016; 21(4):67-76.
53. Otiotoju GTO, Nwamarah JU, Otiotoju O, Odoh EC and Iyeghe LU. Phytochemical composition of some underlisted green leafy vegetables in Nsukka urban L.G.A. of Enugu state. J Biodivers Environ Sci. 2014; 4(4):208-217.
54. Wilfred OO, Philip FU, Emmanuel OE, Sampson C, Nwabuko. Comparative phytochemical, antioxidant and antimicrobial properties of *Ficus capensis*, *Aristolochia ringens*, *Albizia zygia* and *Lannaewelwitschii*. J pharm sci. 2017; 16(2):147-157.
55. Chizoruo IF, Onyekachi IB, Odinaka NP, Chinyelu EM. Phytochemical, FTIR and elemental studies of African Mistltoe (*Viscum album*) leaves on *Cola nitida* from South-Eastern Nigeria. World Scientific News. 2019; 132:84-97.
56. Sampaio BL, Costa FBD. Influence of abiotic environmental factors on the main constituents of the volatile oils of *Tithonia diversifolia*. Rev Bras Farmacogn. 2018; 28(2):135-144.
57. Türkmen, M Kara, M Maral, H. Soylu S. Determination of chemical component of essential oil of Origanum dubium plants grown at different altitudes and antifungal activity against *Sclerotinia sclerotiorum*. J Food Process Preserv. 2022; 46(6):15787.

58. Walia A, Gupta AK, Sharma V. Role of bioactive compounds in human health. *Acta Sci. Med. Sci.* 2019; 3(9):25-33.
59. Lukas S, Jiri M, Lenka S, Mojmir B, Irena H, Tunde J, Rene K, Eva S, Jiri S. Contribution of Red Wine Consumption to Human Health Protection. *Molecule.* 2018; 23(7): 1684.
60. Alugoju P, Periyasamy L, Dyavaiah M. Quercetin enhances stress resistance in *Saccharomyces cerevisiae* tell mutant cells to different stressors. *J Food Sci Technol.* 2018; 55(4):1455-1466.
61. Leri M, Scuto M, Ontario ML, Calabrese V, Calabrese EJ, Bucciantini M, and Stefani M. Healthy effects of plant polyphenols: molecular mechanisms. *Int J Mol Sci.* 2020; 21(4):1250.
62. Ewa N, Irena S, Maciej G, Andrzej M, Piotr S, Joanna P. Małgorzata F. Oxidative Stress in Neurodegenerative Diseases. *Mol Neurobiol.* 2016; 53:4094–4125
63. Pérez-Torres I, Castrejón-Téllez V, Soto ME, Rubio-Ruiz ME, Manzano-Pech L, Guarner-Lans V. Oxidative stress, plant natural antioxidants, and obesity. *Int J Mol Sci.* 2021; 22(4):1786.
64. Medrano-Padial C, Puerto M, Richard T, Cantos-Villar E, Pichardo S. Protection and reversion role of a pure stilbene extract from grapevine shoot and its major compounds against an induced oxidative stress. *J Funct Foods.* 2021; 79:104393.
65. Jesus M, Romaní A, Mata F, Domingues L. Current options in the valorisation of vine pruning residue for the production of biofuels, biopolymers, antioxidants, and bio-composites following the concept of biorefinery: a review. *Polymers.* 2022; 14(9):1640.
66. Naveed M, Hejazi V, Abbas M, Kamboh AA, Khan G J, Shumzaid M, XiaoHui Z. Chlorogenic acid (CGA): A pharmacological review and call for further research. *Biomed Pharmacother.* 2018; 97:67-74.
67. Wang G, Song X, Zhu L, Li Q, Zheng F, Geng X, Sun B. A flavoromics strategy for the differentiation of different types of Baijiu according to the non-volatile organic acids. *Food Chem.* 2022; 374: 131641.
68. Bohn T, Desmarchelier C, El SN, Keijer J, van Schothorst E, Rühl R, Borel P. β -Carotene in the human body: metabolic bioactivation pathways—from digestion to tissue distribution and excretion. *Proc Nutr Soc.* 2019; 78(1):68-87.
69. Mota A, Pinto J, Fartouce I, Correia MJ, Costa R, Carvalho R, Oliveira AA. Chemical profile and antioxidant potential of four table grape (*Vitis vinifera*) cultivars grown in Douro region, Portugal. *Cienc Tec Vitivinic.* 2018; 33(2):125-135.
70. Zhang Z, Zhang J, Fan L, Kilmartin PA. Degradation of cyanidin-3-O-glucoside induced by antioxidant compounds in model Chinese bayberry wine: Kinetic studies and mechanisms. *Food Chem.* 2022; 373:131426.
71. Muhammad Abdul Kadar NN, Ahmad F, Teoh SL, Yahaya MF. Caffeic acid on metabolic syndrome: a review. *Molecules.* 2021; 26(18):5490.
72. Zhu L, Qi X, Bai J, Sun X, Hou H. The mechanism of molecular cross-linking against nonenzymatic degradation in the body wall of ready-to-eat sea cucumber. *Food Chem.* 2022; 373:131359
73. Agunloye OM, Oboh G. Caffeic acid and chlorogenic acid: Evaluation of antioxidant effect and inhibition of key enzymes linked with hypertension. *J Food Biochem.* 2018; 42(4):12541.
74. Dias IH, Milic I, Heiss C, Ademowo OS, Polidori MC, Devitt A, Griffiths HR. Inflammation, lipid (per) oxidation, and redox regulation. *Antioxid Redox Signal.* 2020; 33(3):166-190.
75. Esra AS, Betul AY. Effect of *Taraxacum officinale* L. ethanol extract against kidney injuries induced by paracetamol in rats. *GSC Biol Pharm Sci.* 2022; 21(1):144-151.
76. Odunola OA. Assessment of in-vitro antioxidant activities and genotoxicity in *E. coli* of ethanol extracts of *Vitellaria paradoxa* (Gaertn. F). *Arch Med.* 2019; 7(1):13-20.
77. Tekos F, Makri S, Skaperda Z. V Patouna, A Terizi, K Kyriazis, ID, Demetrios K. Assessment of antioxidant and antimutagenic properties of red and white wine extracts in vitro. *Metabolites.* 2021; 11(7):436
78. Singh S, Gupta A, Verma S. In vitro antioxidant activities of two medicinal plants on the basis of DPPH free radical scavenging activity. *Ann Romanian Soc Cell Biol.* 2021; 13:4807-11.
79. Doris A. Rosemary IU. Phytochemical, Antimicrobial and Free Radical Scavenging Activity of *Ficus capensis* Thunb Leaves. *J Complement Altern Med Res.* 2016; 1(4):1-7
80. Ehimwenma SO, Osarieme TO. Invitro antioxidant activity and phytochemical screening of methanol extracts of *Ficus capensis* and *Dacryodes edulis* leaves. *JPB.* 2014; 11(2):66-75.
81. Ján M, Peter C, Ivana M, Daniel B, Radoslav K. Changes of Selected Antioxidant Parameters of Red Wines during Maturation. *Czech J Food Sci.* 2016; 34 (4):356–361.
82. Gabriele M, Gerardi C, Lucejko JJ, Longo V, Pucci L, Domenici V. Effects of low sulfur dioxide concentrations on bioactive compounds and antioxidant properties of Aglianico red wine. *Food Chem.* 2018; 245:1105-1112.
83. Đorđević NO, Todorović N, Novaković IT, Pezo LL, Pejin B, Maraš gV., and Pajović, S. B. Antioxidant activity of selected polyphenolics in yeast cells: The case study of Montenegrin Merlot wine. *Molecules.* 2018; 23(8):1971